EXHIBIT 38

Nonisotopic DNA Probe Techniques

Nonisotopic DNA Probe Techniques

Edited by

Larry J. Kricka

Department of Pathology and Laboratory Medicine University of Pennsylvania Philadelphia, Pennsylvania



Academic Press, Inc.
Harcourt Brace Jovanovich, Publishers
San Diego New York Boston London Sydney Tokyo Toronto

Front cover photograph: Color enchanced digitized image of a DNA sequence obtained using the chemiluminescent substrate CSPD to visualize bound alkaline phosphatase conjugate. This illustration was kindly provided by Irena Bronstein and Chris Martin of Tropix, Inc.

This book is printed on acid-free paper.

Copyright © 1992 by ACADEMIC PRESS, INC.

All Rights Reserved.

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without permission in writing from the publisher.

Academic Press, Inc.

1250 Sixth Avenue, San Diego, California 92101-4311

United Kingdom Edition published by Academic Press Limited 24–28 Oval Road, London NW1 7DX

Library of Congress Cataloging-in-Publication Data

Nonisotopic DNA probe techniques / edited by Larry J. Kricka.

p. cm.

Includes bibliographical references and index.

ISBN 0-12-426295-3 (hardcover). -- ISBN 0-12-426296-1 (pbk.)

1. DNA probes. I. Kricka, Larry J., date

[DNLM: 1. DNA Probes. 2. Nucleic Acid Hybridization. QU 58

N8124]

QP624.5D73N66 1992

574.87'3282--dc20

DNLM/DLC

for Library of Congress

91-41379 CIP

PRINTED IN THE UNITED STATES OF AMERICA

92 93 94 95 96 97 BC 9 8 7 6 5 4 3 2 1

CONT

Contributor Preface

PART ON: Introduct

1. Nucle Strat

Larry J I.

II. III.

IV.

V. VI.

2. Non

Christo

I.

II. III.

IV.

V. VI. 8

Detection of Horseradish Peroxidase by Colorimetry

Peter C. Verlander
Department of Investigative Dematology
The Rockfeller University
New York, New York

- I. Introduction
 - A. Historical Overview
 - 1. In Situ Hybridizations
 - 2. Sandwich Hybridizations
 - 3. Membrane Hybridizations
 - B. Substrates
 - 1. Insoluble Products
 - 2. Soluble Products
- II. Materials
- III. Procedures
 - A. In Situ Hybridizations
 - 1. AEC Detection
 - 2. DAB Detection
 - B. Microtiter Plate Assays
 - 1. TMB Detection
 - 2. OPD Detection
 - C. Membrane Hybridizations
 - 1. AEC Detection
 - 2. TMB Detection
- IV. Conclusions
 - References

I. INTRODUCTION

A. Historical Overview

Horseradish peroxidase (HRP) has been used extensively as a colorimetric marker in biological studies. A hemoprotein with a molecular weight of 40,000, HRP is an ideal detection reagent because of its stability, high turnover rate, and the availability of a wide assortment of colorimetric substrates. Although radioactive techniques are generally more sensitive, the sensitivity of peroxidase detection systems is adequate for many applications, and speed and safety considerations make peroxidase detection

). Verlander

dapted to / derived I method 1 (Langer f a biotisequential e, or by a tavidin or ptavidin—ed of peritive than

oxidase to nunologic nodeoxyu-1989). Imof biotioupled to ted into a sandwich

been most igati et al., rapid than than days, sensitivity ly involves ne ultimate lex. Sensi, have been per cell are marker for 25, and has




Fig. 1 Detection of nucleic acid probes with horseradish peroxidase. (A) Hybridization with a biotinylated probe, followed by sequential washes with streptavidin and biotinylated peroxidase. (B) Hybridization with a biotinylated probe, followed by a single wash with a preformed complex of streptavidin and biotinylated peroxidase. (C) Hybridization with a biotinylated probe, followed by detection with antibiotin antibodies coupled to peroxidase. This technique also applies to detection of probes modified in other ways, such as by sulfonation. (D) Hybridization with a probe that is directly coupled to peroxidase.

been used for a variety of clinical diagnostic tests. Peroxidase detection has also been used for genetic testing applications, such as Y chromosome aneuploidy (Guttenbach and Schmid, 1990).

2. Sandwich Hybridizations

Sandwich assays are particularly well suited to peroxidase detection. These assays generally include a capture probe that is bound to a fixed matrix, such as nitrocellulose membranes (Dunn and Hassell, 1977; Ranki et al., 1983), microtiter wells (Dahlen et al., 1987; Keller et al., 1989), or

188 Peter C. Verlander

beads (Langdale and Malcolm, 1985; Polsky-Cynkin et al., 1985). The target nucleic acid molecule hybridizes to the capture probe and is thereby bound to the matrix, while a second probe that is either directly or indirectly labeled with peroxidase hybridizes to an adjacent sequence on the target (Fig. 2). These techniques are particularly useful for the detection of polymerase chain reaction (PCR)-amplified nucleic acids, and have been used in assays for human immunodeficiency virus (HIV) (Kemp et al., 1990; Keller et al., 1989), β -thalassemia (Saiki et al., 1988, 1989), and sickle-cell anemia (Saiki et al., 1988) among others. Peroxidase detection is useful for these types of assays because of the high turnover rate of the enzyme, and because of the availability of a number of sensitive substrates for soluble assays.

3. Membrane Hybridizations

Peroxidase detection is perhaps less well suited to membrane hybridization techniques. The sensitivity of peroxidase detection does not match that of radioactive systems, although detection of subpicogram bands on Southern hybridizations have been claimed (Sheldon *et al.*, 1986, 1987). The lower stability of peroxidase as compared to alkaline phosphatase




Fig. 2 Sandwich assays. The target hybridizes to an immobilized probe and is captured to the solid matrix. A biotinylated probe hybridizes to adjacent sequences on the target and is detected by streptavidin-HRP.

8. Horser.

limits t nonspe tion ha 1988, l' membr histocc

B. Su

A wide ating b marily but oth and the ations.

1. Ir

The situ h ethylca naphth of their describ table a less co Initia benzid a tende of gen 1964). organic hybrid photog suspec

of the ta very ity of Soban substrate by the

DAF